BLACK GRAM BEAN

Urd, Urd Bean, Black Gram

Vigna mungo

Fabaceae

ECHO® PLANT INFORMATION SHEET

Origin

This very important bean in the diet of vegetarian Indians was originally found growing in India. It is now widely cultivated in the tropics.

Uses

Black Gram is perhaps the most important pulse crop in Asia because of its versatility of use. The bean is used whole, split, husked, fermented, sprouted or ground into flour which when combined with rice flour is especially valued for baking. After the plants are picked, animals can graze it and it can be plowed under as green manure. It is a legume with a deep root system that binds soil particles together preventing erosion. Its foliage is lush so that when cut and left on the ground, it acts as good mulch.

Common Names

• French

haricot mungo

Cultivation

Black Gram is an annual, warm season crop, which produces pods of 4-6 cm (1.5" - 2.5 in) in 3 months. It tolerates a wide range of soils often succeeding in heavier soils where other crops will not. It can be planted in the late rainy season after maize or soybeans. If the field has previously grown a legume, black gram will freely form nitrogen-fixing nodules on its roots.

Harvesting and Seed Production

The pods turn black when mature though not all will mature at once so 2-4 hand pickings at 10-15 day intervals may be necessary. Plants are often pulled up as the pods mature, left in the field to dry for a week then threshed either by beating with a stick or trod by bullocks. The seeds may be further dried in the sun and stored in a dry area away from rodents.

Pests and Diseases

The threat of various fungi can be overcome by growing black gram in irrigated paddy fields rather than planting the crop in the rainy season.

Cooking and Nutrition

Immature pods of the Black Gram bean can be eaten as a cooked vegetable. Soaked beans can be sprouted or combined with vegetables and spices to make dal, fried crisp as papads or in a fermented batter called idli. The bean is richer in protein than most other grains.

References

http://ecocrop.fao.org/ecocrop/srv/en/cropView?id=2149