

GALLITO (AGATÍ DE LA INDIA, COBREQUE, CRESTA DE GALLO, PICO DE FLAMINGO)

Sesbania grandiflora

ECHO® FICHA DE INFORMACIÓN SOBRE PLANTAS

Descripción

Sesbania is a small, quick-growing, loosely-branching tree with compound, alternate leaves. Large pea-like flowers, white to pink, grow in clusters of 2-5 at the base of leaves. Bears narrow pods 30-50 cm long and 8 mm wide.

Usos

Crop uses (culinary) – *Sesbania* flowers are eaten both cooked and raw, in salads, curries, and as a steamed vegetable. Young leaves and green pods can also be cooked and eaten. The dried leaves are used in some countries as a tea, which is considered to have antibiotic, anti-helminthic, anti-tumour, and contraceptive properties.

Crop uses (soil improvement) – *Sesbania* is a nitrogen fixing species which can be used as a green manure, either planted in the field as an annual cover crop or grown on rice paddy bunds. It has been reported to cover the ground in less than four weeks, and grow to a height of 6 feet (1.8 m) in 50 days, producing over 2 tons per acre (about 4.5 kg per hectare) of biomass (Nolte). When grown at the edges of fields or on paddy bunds, the side branches can be trimmed and incorporated as a green manure. It also casts sparse shade, making it a suitable inter-cropping species.

Crop uses (livestock production) – *Sesbania* is a high-quality livestock feed. It contains over 30% crude protein, with low levels of condensed tannins and high digestibility. Due to the presence of some anti-nutritive substances, however, *sesbania* should not be fed to chickens or other non-ruminants. Some authors recommend limiting ruminant's consumption of *sesbania* to 30% of their diet, although studies in Indonesia report up to 70% consumption in the dry season, when other species are not available.

Other uses – *Sesbania* can be used to cast light shade for tree nurseries or vegetable crops, as a living trellis for vining plants, and as a windbreak. *Sesbania* wood can be used for pulp and paper production.

Cultivo

Seasons of production – In Thailand, *Sesbania* flowers in the hot season, around April-May. As a fodder, it can be harvested for most of the year, at 3-4 month intervals.

Length of production and harvest period – When managed for fodder, side branches are removed every 3-4 months. Repeated pruning of main stem shortens the life of the tree. Some farmers use *sesbania* as an annual cover crop, planting into rice paddies then harvesting or tilling in the foliage before the next cropping cycle.

Pollination – The large hermaphroditic flowers are pollinated by birds.

Plant spacing – Commonly planted as individual trees or in rows, space 1-2 m (2 - 6.5 ft) apart in fence lines, field borders and on the bunds of rice paddies. Can grow as much as 5-6 m (16-20 ft) in its first 9 months. Can be planted in high densities (up to 3000 plants/ha) for pole timber production, or in low densities to produce dry season forage and fuelwood. For reforestation, it is recommended to space seedlings or 1 m long cuttings at 4x4 m (13x13 ft).

Production methods – Proper cutting management can greatly influence the productivity of *Sesbania*, as it cannot survive repeated cuttings. Farmers in Lombok, Indonesia cut only the side branches for fodder, leaving the main growing stem untouched. When foliage is no longer within reach, the trees are felled. The long straight poles can be used for firewood or light construction (although it is reported that *Sesbania* firewood burns poorly and produces much smoke). *Sesbania* has been grown in windbreaks and in association with guinea grass. The sparse canopy and erect habit enables it to be grown on crop margins with little reduction in sunlight, while still providing N fixation. It can also be used to shade nurseries and gardens. The harvesting of side branches has yielded 4.5-9 kg (10-20 lb) of leaves per year. A green manure yield of 55 t/ha of green material in 6.5 months was achieved in Java. Wood yields of 20-25 m³/ha/year have been achieved in commercial plantations in Indonesia.

Known environmental conditions for production – *Sesbania* is well adapted to hot, humid environments and does not grow well in the subtropics with cool temperatures reaching below 10°C (50° F). It is reportedly outstanding in its ability to tolerate waterlogging, which makes it ideally suited to seasonally waterlogged or flooded environments. Is adapted to rainfall conditions of 200 – 400 cm (80 – 160 in) but will grow in areas receiving only 80 cm (31 in).

Known soil requirements – *Sesbania* is known to tolerate both saline and alkaline soil conditions. It grows best in soils with pH 6.6 to 8.5, but is tolerant of a high variety of soil types, including heavy and very poor soils.

La cosecha y producción de semillas

Seed saving – *Sesbania* can be propagated by cuttings as well as seed. It seeds in lengthening days in pods of 15-50 seeds. Allow pods to dry before

breaking open, then dry seeds and store. Viability deteriorates more rapidly than most seeds. Sesbania seed germinates easily without scarification, although scarification has been shown to increase evenness of germination.

Plagas y Enfermedades

Known pests – Sesbania is thought to be relatively pest free in some environments, but it is susceptible to severe attacks from leaf webbers (*Azygophleps scalaris*), leaf feeders and stem borers, and can be highly susceptible to root-knot nematode (*Melodogyne incognita*). It is also susceptible to grey leaf spot (*Pseudocercospora sesaniae*), and Sesbania mosaic virus is reported in India.

Referencias

James A. Duke. 1983. Handbook of Energy Crops. unpublished.

"Sesbania grandiflora (L.) Pers.." Center for New Crops and Plants Products. Purdue University. Web. http://www.hort.purdue.edu/newcrop/duke_energy/Sesbania_grandiflora.html

"Sesbania grandiflora." Tropical Forages. FAO Collaboration. Web. http://www.tropicalforages.info/key/Forages/Media/Html/Sesbania_grandiflora.htm' Suttie, J.M..

"Sesbania grandiflora (L.) Poir." CIAT/FAO collaboration on Tropical Forages. FAO. Web. <http://www.fao.org/ag/AGP/AGPC/doc/Gbase/Default.htm>