

# CANISTEL

*Pouteria campechiana*

Sapotaceae

ECHO® KARATASI YA MAELEZO YA KUPANDA

## Maelezo

The Canistel is a large, evergreen tree native to Central America. It usually grows to around 8 m in height, but occasionally grows larger in favorable conditions. Leaves are commonly oblanceolate at 10-30cm in length and alternately cluster around branch tips. The tree has an open-growing canopy, light brown mature bark, and abundant latex. The fruit shape and size is variable. Green, immature fruits ripen to a golden-yellow or orange-yellow. The fruit varies from firm and fibrous towards the exterior, to a cooked egg yolk texture towards the center. The flavor is sweet and mild, often described as a cooked sweet potato.

## Majina ya Kawaida

- English
  - Canistel
  - Egg Fruit
  - Yellow Sapote

## uvunaji na uzalishaji wa mbegu

Mature fruits should be allowed to ripen fully before consumption to reduce the quantity of latex. Yellow fruits will ripen on the tree, but they are commonly picked when fully yellow and allowed to soften off the tree (3-10 days after harvest). Fruits should be clipped off trees to avoid damaging the fruit. The seeds should be planted soon after harvest, and may not germinate for several months (3-5 months average). Seeds should remain in well-drained, damp composted soil throughout the germination period.

## wadudu na magonjwa

The trees are often very healthy and are not usually attacked by pests. Scale insects are the most common pest, with rust and fruit spot occasionally damaging fruit .

## Marejeleo

Morton, J. 1987. Canistel. p. 402-405. In: Fruits of warm climates. Julia F. Morton, Miami, FL.

<http://ecocrop.fao.org/ecocrop/srv/en/cropView?id=2394>